



Clean Genome® *E. coli*
MDS™ 42 Pack, Electrocompetent Cells
(Cat. No. E-1000-15)

FOR RESEARCH USE ONLY

COMPONENTS

E-1000-15

MDS™ 42 Electrocompetent Cells	0.2 ml
MDS™ 42 <i>recA</i> ⁻ Electrocompetent Cells	0.2 ml
MDS™ 42 <i>recA</i> ⁻ Blue Electrocompetent Cells	0.2 ml
pUC19 Control DNA (10 pg/μl)	50 μl
SOC Medium	2 x 10 ml

STORAGE CONDITIONS

Store components at -80°C. **Do not store cells in liquid nitrogen.**

BACKGROUND

Using synthetic biology methods, the *Escherichia coli* K-12 genome was reduced by making a series of planned, precise deletions. The multiple-deletion series (MDS™) strains (1), with genome reduction of up to 15%, were designed by identifying non-essential genes and sequences for elimination, including recombinogenic or mobile DNA and cryptic virulence genes, while preserving robust growth and protein production. Genome reduction also led to unanticipated beneficial properties, including high electroporation efficiency and accurate propagation of recombinant genes and plasmids that are unstable in other strains. Subsequent deletions and introduction of useful alleles produce strains suitable for many molecular biology applications.

GENOTYPES

MG1655 multiple-deletion strain (1)
MG1655 multiple-deletion strain(1), *recA*⁻ 1819
MG1655 multiple-deletion strain(1), *recA*⁻ 1819, *lacZ* M15⁻

The *recA* 1819 mutation is a complete deletion of *recA*⁻.

The *lacZ* M15 deletion has been created in the genome to allow blue/white screening of inserts in plasmids using the α-complementing fragment of β-galactosidase.

ELECTROCOMPETENT CELL QUALITY CONTROL

Transformation efficiency is tested using pUC19 Control DNA, in duplicate. Transformed cells are plated onto LB plates containing 50 μg/ml carbenicillin. Transformation efficiency is > 5 x 10⁹ cfu/μg DNA.

BEFORE YOU BEGIN

Please note, Clean Genome® strains do not remain viable for extended periods (greater than 2 weeks) when stored at 4°C. We recommend preparing glycerol stock cultures of clones and storing at -80°C, or keeping plates at room temperature for up to 5 days.

To ensure that the cells grow on minimal media and to prevent a significant lag when transferring to liquid culture, we streak from glycerol stocks of clones onto minimal plates with 0.2% glucose and grow at 37°C for 24 h, at 30°C for 48 h, or at room temperature (RT) for several days (e.g., over the weekend). Colonies picked from these plates are used for cultures. If you are only using rich media (e.g., LB, TB) for all growth, rich medium plates can be used for streaking clones.

Clean Genome strains do not have flagella and tend to aggregate and drop fairly quickly from solution. To obtain OD readings, cells should be mixed just before taking an aliquot for dilution, and dilution samples should be mixed just before taking an OD reading.

PRE-TRANSFORMATION STEPS

1. Thaw the provided SOC medium and warm it to 37°C to dissolve any visible precipitate.
2. Warm selective antibiotic plates to room temperature or 37°C.
3. If you are transforming a plasmid with antibiotic resistance other than ampicillin, you will need at least one LB agar plate containing 100 µg/ml ampicillin (or 50 µg/ml carbenicillin) for plating the pUC19 Control DNA transformation.
4. If you are planning to use MDS™ 42*recA*⁻ Blue for blue/white screening, you will need LB agar plates containing the appropriate antibiotic, 80 µg/ml 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal), and 0.5 mM Isopropyl-β-D-thiogalactopyranoside (IPTG). Alternatively, spread 20 µl 50 mg/ml X-Gal (in dimethylformamide) and 100 µl 0.1 M IPTG onto LB agar plates containing the appropriate antibiotic and incubate at least 1 h at 37°C, so that the plates absorb the X-Gal and IPTG before plating.

TRANSFORMATION PROCEDURE

1. Place the required number of **0.1-cm** electrode gap electroporation cuvettes and 1.5-ml microcentrifuge tubes on ice.
2. Add 1-2 µl experimental DNA to 1.5-ml microcentrifuge tubes.

NOTE - Transformation efficiency of ligation products can be increased several fold by diluting the ligation reaction 5-fold with TE or water prior to adding DNA.

3. For the positive control, add 1 µl of pUC19 Control DNA to a microcentrifuge tube. For the negative control, add 1 µl Ultrapure H₂O.
4. Thaw Clean Genome MDS™ 42 Electrocompetent Cells on ice and use as soon as possible.
5. When the cells are thawed, flick the tube gently 1-3 times to evenly suspend cells. Add 40 µl cells to each pre-chilled microcentrifuge tube containing DNA.
6. Gently pipet the cell/DNA mixture 1-2 times to mix. Discard any unused cells. **Re-freezing cells is not recommended.**
7. Pipette the cell/DNA mixture into a pre-chilled 0.1-cm electroporation cuvette avoiding bubble formation. Gently tap the cuvette several times to remove bubbles and to ensure the cell/DNA mixture is at the bottom of the cuvette.
8. Electroporate samples at 18 kV/cm. Add 960 µl of room-temperature SOC medium to the cuvette of electroporated cells. Mix well by pipetting up and down 2-3 times. Transfer the mixture to a 15-ml culture tube.
9. Incubate at 37°C with shaking at 250–275 rpm for 1 h.
10. For cells transformed with pUC19 Control DNA, dilute the culture 1/50 in SOC medium and plate 100 µl of the dilution onto pre-warmed LB plates containing 100 µg/ml ampicillin.
11. For cells transformed with experimental DNA, plate as required and spread up to 100 µl onto selective plates. **OPTIONAL:** For blue/white screening, spread cells onto LB plates containing the appropriate antibiotic, 80 µg/ml X-Gal, and 0.5 mM IPTG.
12. Incubate plates overnight at 37°C.

REFERENCE

1. Pósfai, G., Plunkett III, G., Fehér, T., Frisch, D., Keil, G.M., Umenhoffer, K., Kolisnychenko, V., Stahl, B., Sharma, S.S, de Arruda, M., Burland, V., Harcum, S.W., and Blattner, F.R. Emergent Properties of Reduced-Genome *Escherichia coli*. *Science* **312**, 1044-1046 (2006).

CALCULATION OF TRANSFORMATION EFFICIENCY USING pUC19 CONTROL DNA

1. Count cells on each Control plate. Calculate the average colonies/plate for the pUC19 Control DNA plates if more than one replicate was plated (Note: the No DNA Control plate should not have any colonies).
2. Use the formula below to calculate the transformation efficiency in colony forming units (cfu) per µg plasmid Control DNA.

$$\frac{\text{Average \# colonies}}{\text{pg DNA plated}} \times \frac{10^6 \text{ pg}}{\mu\text{g}} = \frac{\text{cfu}}{\mu\text{g plasmid DNA}}$$

EXAMPLE OF TRANSFORMATION EFFICIENCY USING pUC19 CONTROL DNA

$$\frac{10 \text{ pg DNA}}{1000 \mu\text{l total rxn vol.}} \times \frac{1}{50} \text{ dilution} \times 100 \mu\text{l plated} = 0.02 \text{ pg DNA plated}$$

$$\frac{200 \text{ colonies}}{0.02 \text{ pg DNA}} \times \frac{10^6 \text{ pg}}{\mu\text{g}} = 1.0 \times 10^{10} \frac{\text{cfu}}{\mu\text{g}}$$

TROUBLESHOOTING

The pUC19 Control DNA plasmid provided should be used as a positive control.

Problem	Possible Solution
Arcing or “popping” occurs during electroporation	1. DNA may contain salt. Desalt DNA sample or dilute it 1/5 before electroporation. Repeat electroporations with 1 µl of the desalted or diluted DNA.
No colonies or low number of colonies	1. Incorrect drug selection or drug concentration. Verify that LB agar plates contained the appropriate selective antibiotic. Repeat electroporations and plate on new plates. 2. Plates were too old/dry. Repeat electroporations and plate on new plates. 3. Cells were not handled correctly. Cells must be handled very gently. Do not pipet vigorously or vortex. Always gently pipet or gently flick cells to resuspend. Thaw cells and keep on ice until electroporation. Repeat electroporation.
Lawn or confluent cell growth or satellite colonies	1. Incorrect drug concentration. Verify that LB agar plates contain the appropriate selective antibiotic concentration. Repeat electroporation and plate onto new plates. 2. Antibiotic has degraded. Plates are too old, antibiotic stock(s) have degraded, or antibiotic was added when medium was too hot. Prepare fresh antibiotic stock(s) and fresh plates. Repeat electroporation. 3. Incubated at 37°C too long. Plates should incubate for 16-18 h. Amp ^R cells secrete β-lactamase that creates a drug-free zone in the surrounding medium, allowing small amp ^S colonies to grow. Carbenicillin (an ampicillin analog) appears to be less vulnerable to degradation. Repeat Electroporation and incubate plates for 16-18 h.
Colonies from pUC19 control DNA plates are white or light blue when plated on LB agar plates containing X-Gal and IPTG.	1. Colonies have not grown long enough or are very small due to high plating density. Extend the 37°C incubation or keep the plates at RT, to allow time for better expression of the β-galactosidase enzyme and color development. The best color development occurs 24 h after plating. 2. Colonies are too small due to old/dry plates. Repeat electroporation and plate on new plates. 3. Incorrect X-Gal and/or IPTG concentration in the plates. Verify that LB agar plates contain the appropriate X-Gal and IPTG concentrations. Repeat electroporation.

LIMITED PRODUCT WARRANTY

Recipient acknowledges that the Material is experimental and is supplied to Recipient WITHOUT ANY WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF TITLE, MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY WARRANTIES REGARDING INFRINGEMENT OF THIRD PARTY RIGHTS. Recipient agrees to rely solely upon its own opinion of the Material with regard to their safety and suitability for any purpose.

Recipient agrees to waive all claims against Scarab except as may be limited by state law governing Recipient. Recipient agrees to defend and indemnify Scarab, its employees and agents from all claims asserted by any third party and any damages and recoveries arising from the use, storage, or handling of the Material caused by or allowed by Recipient. Scarab makes no representation that the use of the Material will not infringe any patent or proprietary rights of any third parties.

USAGE RESTRICTIONS (LICENSING)

Scarab is providing you with this Material subject to the non-transferable right to use the subject amount of the Material for your research at your academic institution. The Recipient agrees not to sell or otherwise transfer this Material, or anything derived or produced from the Material to a third party. NO RIGHTS ARE PROVIDED TO USE THE MATERIAL OR ANYTHING DERIVED OR PRODUCED FROM THE MATERIAL FOR COMMERCIAL PURPOSES. If the Recipient makes any changes to the chromosome of the Material that results in an invention in breach of this limited license, then Scarab will have a worldwide, exclusive, royalty-free license to such invention whether patentable or not.

If the Recipient is not willing to accept the terms of this limited license, Scarab is willing to accept return of this product with a full refund, minus shipping and handling costs. For information on obtaining a license to this Material for purposes other than research, please contact Scarab's Licensing Department.

Clean Genome[®] is a registered trademark of Scarab Genomics, LLC.

Scarab Genomics' technology is covered by U.S. Pat. No. 6,989,265 and related foreign applications.

© 2012, Scarab Genomics LLC. All rights reserved.